

()

2017 :

:

04 :

:

(04):

.10 3^n n - (1

$A_n = 3^{16n+6} - 2 \times 109^{2n+3} - 13$: 10 A_n -

$(3n+4) \times 9^n + 7^{2n+1} \equiv 3^{2n} (3n+1) [10] : n$ - (2

.10 $(3n+4) \times 9^n + 7^{2n+1}$ n -

$\overline{y611}$ 3 $\overline{xx0xx01}$ A (3

.7

. A y x -

10 3^n 4 (4


.2017

X -

X

(04):

OABCDEFG



$(O; \overrightarrow{OA}, \overrightarrow{OC}, \overrightarrow{OD})$

(ACD) $\vec{n}(1,1,1)$ - (1

(ACD)

(ACD) O (Δ) (2

(Δ)

(ACD) (Δ) H -

$$M(x, y, z) \quad (S_m) \quad (3)$$

$$x^2 + y^2 + z^2 - 2mx - 2my - 2mz - 1 + 3m^2 = 0 \quad (m \in \mathbb{R})$$

$$(S_m) \quad m \quad -$$

$$(S_m) \quad A \quad m \quad -$$

$$(\Delta) \quad \begin{pmatrix} S_2 \\ S_3 \end{pmatrix} (S_0) \quad \omega_{\frac{2}{3}} \quad \omega_0 \quad - \quad (4)$$

$$\begin{pmatrix} S_2 \\ S_3 \end{pmatrix} (S_0) \quad (ACD) \quad -$$

:(05): _____

$$E \quad D \quad C \quad B \quad A \quad (O; \vec{u}; \vec{v})$$

$$z_E = -2i \quad z_D = -1+i \quad z_C = 3i \quad z_B = 4+i \quad z_A = 1$$

$$C \quad B \quad E \quad D \quad T \quad \frac{z_C - z_A}{z_B - z_A} = \frac{z_E - z_A}{z_D - z_A} \quad (1)$$

$$\frac{\sqrt{2}}{2} \quad \frac{\pi}{4} \quad A \quad S \quad C \quad C' \quad (2)$$

$$[EB] \quad [DE] \quad [CD] \quad [BC] \quad I_4 \quad I_3 \quad I_2 \quad I_1 \quad (3)$$

$$I_2 \quad I_4 \quad I_1 \quad r \quad -$$

$$I_1 \quad I_2 \quad I_3 \quad I_4 \quad z_{I_2} + z_{I_4} \quad z_{I_1} + z_{I_3} \quad -$$

$$S \quad z' \quad M' \quad z \quad M \quad (4)$$

$$z' = \frac{1}{2} [(1+i)z + 1-i] \quad *$$

$$\theta \in \left[0; \frac{\pi}{2}\right] \quad z = (i-1)(1+e^{i\theta}) \quad Z \quad M \quad (\gamma) \quad (5)$$

$$\left[0; \frac{\pi}{2}\right] \quad \theta \quad (\gamma) \quad -$$

أوجد طبيعة المجموعة (γ') صورة (γ) -

:(07): _____

$$f(x) = (3+x)e^{\frac{-x}{2}} \quad : \quad \mathbb{R} \quad f$$

$$(2cm :) \quad (O; \vec{i}; \vec{j}) \quad (C_r)$$

$$. +\infty \quad -\infty \quad f \quad - \quad (1)$$

$$. \mathbb{R} \quad f \quad -$$

$$. -2 < \alpha < \frac{-3}{2} : \quad \alpha \quad \mathbb{R} \quad f(x) = 3 \quad - \quad (2)$$

$$. (O; \vec{i}; \vec{j}) \quad (C_f) \quad -$$

$$. \mathbb{R} \quad f(x) = m \quad m \quad . \quad m \quad -$$

$$I = \int_{-3}^0 x e^{-\frac{x}{2}} dx \quad (3)$$

$$. -3 \leq x \leq 0 \quad 0 \leq y \leq f(x) : \quad M(x; y)$$

$$. g(x) = 3e^{\frac{x}{2}} - 3 : \quad \mathbb{R} \quad g \quad (4)$$

$$. g(x) = x \quad f(x) = 3 \quad -$$

$$. (g \quad g') . \mathbb{R} \quad g \quad g' \quad -$$

$$. g'(\alpha) = \frac{\alpha + 3}{2} : \quad -$$

$$: [-2; \alpha] \quad x \quad (5)$$

$$. [-2; \alpha] \quad g(x) -$$

$$. \frac{1}{2} \leq g'(x) \leq \frac{3}{4} -$$

$$. 0 \leq \frac{1}{2}(\alpha - x) \leq g(\alpha) - g(x) \leq \frac{3}{4}(\alpha - x) : [-2; \alpha] \quad x \quad (6)$$

$$. u_{n+1} = g(u_n) : n \quad u_0 = -2 : \quad \mathbb{N} \quad (u_n) \quad (7)$$

$$. -2 \leq u_n \leq \alpha : n \quad -$$

$$. 0 \leq \alpha - u_n \leq \left(\frac{3}{4}\right)^n \quad 0 \leq \alpha - u_{n+1} \leq \frac{3}{4}(\alpha - u_n) : n \quad -$$

$$. u_n \quad -$$

: (04):

- .7 $a = p^4 - 1$ p a
- .3 a 3 (-1) 1 p (1)
- .16 a $p^2 - 1 = 4k(k+1)$: k (2)
- . $a \equiv 0 [5]$ 5 p (3)
- . δ β α (4)
- . δ $\alpha\beta$ β α δ β δ α -
- . a 240 -

: (04):

- الفضاء منسوب إلى المعلم المتعامد والمتجانس $(O; \vec{i}, \vec{j}, \vec{k})$:
- $E(2;3;-1)$ $D(1;3;1)$ $C(4;4;1)$ $B\left(4 - \frac{3\sqrt{2}}{2}; 4 + \frac{3\sqrt{2}}{2}; 4\right)$ $A\left(4 + \frac{3\sqrt{2}}{2}; 4 - \frac{3\sqrt{2}}{2}; 4\right)$
- . $C; B; A$ (1)
- . (ABC) D (Δ) - (2)
- . (Δ) E -
- . $[AB]$ Ω (3)
- . (S) $C; B; A$ -
- . m $x + y - mz + 4m - 8 = 0$ (P_m) - (4)
- . (d)
- . $z = 4$ (Q) (P_0) -
- . (Q) (S) (S') -

: (05):

- . $\frac{3\pi}{4}$ $\frac{1}{2}$ A_0 S $A_0 B_0 = 8$ B_0 A_0
- . $B_{n+1} = S(B_n)$ n : (B_n)
- . B_3 B_2 B_1 (1)
- . $A_0 B_{n+1} B_{n+2}$ $A_0 B_n B_{n+1}$ n (2)
- . $(\overline{A_0 B_0}, \overline{A_0 B_n}) \equiv \frac{3\pi}{4} n [2\pi] : n$ (3)

$$u_n = B_n B_{n+1} \quad n \quad : \quad (u_n) \quad (4)$$

$$u_0 \quad n \quad u_n \quad q \quad (u_n) \quad -$$

$$\lim_{n \rightarrow +\infty} T_n \quad T_n = u_0 + u_1 + u_2 + \dots + u_n \quad n \quad -$$

$$3x - 4y = 2 : \quad \mathbb{Z} \times \mathbb{Z} \quad - \quad (5)$$

$$n \quad A_0 \quad (A_0 B_0) \quad (\Delta) \quad -$$

$$(\Delta) \quad B_n$$

:(07):

$$g(x) = \ln\left(1 + \frac{1}{x}\right) - \frac{1}{1+x} : \quad]-\infty; -1[\cup]0; +\infty[\quad g \quad (1)$$

$$]-\infty; -1[\cup]0; +\infty[\quad g(x) \quad g \quad -$$

$$: \quad D =]-\infty; -1[\cup]0; +\infty[\quad f \quad (2)$$

$$\begin{cases} f(x) = x \ln\left(1 + \frac{1}{x}\right); x \in]-\infty; -1[\cup]0; +\infty[\\ f(0) = 0 \end{cases}$$

$$0 \quad f \quad -$$

$$\lim_{|x| \rightarrow +\infty} f(x) = 1 \quad -$$

$$f \quad f'(x) = g(x) : \quad]-\infty; -1[\cup]0; +\infty[\quad x \quad -$$

$$(2cm) \quad (o; \bar{i}, \bar{j}) \quad f \quad (C_f) \quad (3)$$

$$h(x) = f(-1-x) : \quad h \quad (4)$$

$$]-\infty; -1[\cup]0; +\infty[\quad h \quad -$$

$$(\quad) \quad h \quad -$$

$$x = -\frac{1}{2} \quad (C_f) \quad h \quad (C_h) \quad -$$

$$(C_h) \quad (5)$$

$$\int_{\frac{1}{2}}^1 [1 - f(x)] dx \quad (6)$$

